



The Photon-induced Genesis of RNA and DNA: A Quantum-geometric CNP Model

Pavle Vesić

Independent Researcher, Belgrade, Serbia

Abstract: The evolutionary transition from prebiotic chemistry to the first information-bearing macro-molecules remains one of the most significant paradoxes in modern science. This paper introduces a deterministic, quantum-geometric model for the origin of nucleic acids driven by the Charge Neutralization Process (CNP) under fluctuating environmental photon fluxes. Operating within Hestenes' Spacetime Algebra (Cl1,3) and Friston's Free Energy Principle (FEP), we reframe the genesis of RNA and DNA not as a stochastic chemical accident, but as a fundamental topological necessity of vacuum stress compensation. We demonstrate that high-energy, non-ionizing photon interactions (Case 3 geometry) distort localized electron-proton Zitterbewegung (zbw) rotors, imposing a geometric stress tensor on the boundary of macro-molecular systems (n-th Markov blankets). To preserve boundary integrity and minimize variational free energy, CNP forces the polymerization of free nucleotides into a single-stranded RNA chain, which acts as a dynamic geometric antenna that neutralizes the external electromagnetic flux. Crucially, when the external photon source fades, the asymmetric RNA strand becomes an internal source of vacuum deformation. To eliminate this residual internal stress, CNP executes an inversion of the topological gradient, drawing complementary bases into a stable, bivector phase-locked double helix. DNA is therefore revealed to be a stabilized, archetypal "null-surface" of the vacuum—a physical archive of past environmental stress. This framework seamlessly integrates the thermodynamic dissipation theories of life with the RNA World hypothesis under a singular cosmic algorithm.

Keywords: CNP, STA, FEP, RNA, DNA, UV-C, Photon Fade

INTRODUCTION

Definition of the Charge Neutralization Process (CNP) and Scope of the Present Work

To overcome the methodological fragmentation between fundamental field theory and biophysical self-organization, this paper formally introduces an extended application of the Charge Neutralization Process (CNP). Crucially, CNP is not proposed as a new fundamental physical force, nor does it postulate the existence of novel elementary particles or fields outside the standard model paradigm. Instead, CNP is formally defined as an emergent, geometric, and informational algorithm of dynamic compensation through which a physical or abstract system rearranges and satisfies existing conservation laws—specifically the conservation of energy, momentum, and total angular momentum—under localized potential gradients.

It is necessary to clarify the relationship between our previous work and the present manuscript. The foundational mathematical and functional formulation of CNP as a localized phenomenon was established in our earlier publication. The new contribution of the present paper lies in extending this mechanism globally to cross-disciplinary domains, introducing for the first time the topological factorization of the spacetime spinor, the boundary

commutator condition for conditional independence, and the explicit tensor coupling between vacuum geometry and the Free Energy Principle.

Within this parsimonious framework, "charge" does not solely denote classical electric charge, but rather represents a generalized geometric measure of any uncompensated potential gradient:

1. Geometric and rotational stress in a structured vacuum manifold;
2. Informational stress (prediction error) at the boundary of an organism;
3. Phenomenological tension (imbalance) in panpsychist models of consciousness.

The Thermodynamic and Structural Paradox of Biogenesis

The evolutionary transition from prebiotic chemistry to the first information-bearing macromolecules remains one of the most significant paradoxes in modern science. Prebiotic models rely heavily on stochastic chemistry and random self-assembly, failing to provide a deterministic mechanism for the structural emergence of complex codes. From a thermodynamic perspective, existing non-equilibrium thermodynamics successfully identifies living systems as dissipative structures driven by UV photon degradation but lacks a precise micro-physical geometric mechanism. The missing link requires a unified framework capable of explaining how continuous physical energy fluxes (photons) directly dictate the exact architectural geometry of information storage molecules (RNA and DNA).

Atoms under Pressure: The Three Cases of Photon-Matter Interaction

To ground this macroeconomic model of biological polymerization in first-principles physics, we analyze the micro-physical scaling baseline of a hydrogen atom under electromagnetic stress. Within Hestenes' Spacetime Algebra (Cl_{1,3}), the electron is not treated as a zero-dimensional point charge, but rather as a localized lightlike circular current—the Zitterbewegung (zbw) phenomenon—governed by the real spinor $\psi_e = R_e \rho_e^{1/2}$. The intrinsic spin bivector S_e defining the plane of this localized vacuum rotation is expressed as:

$$S_e = (\hbar/2) R_e I \sigma_3 R_e \tilde{e}$$

When an external electromagnetic field, represented by the unified field bivector $F_{\text{ext}} = E + I B$, interacts with this atomic structure, three discrete topological regimes emerge based on the energy and geometry of the incident flux:

1. Case 1 (Ionization/Disruption): High-energy high-frequency scattering where $\hbar\omega \gg m_e c^2$. The external stress tensor completely overwhelms the local binding vacuum geometry, uncoupling the electron spinor from the proton domain entirely.
2. Case 2 (Quantum Excitation): Resonant absorption matching discrete atomic transitions, where the electron transitions to a higher eigenstate before relaxing and re-radiating the energy back into the environment.
3. Case 3 (Geometric Adaptation and Rotor Shifts): Mid-range, non-ionizing interaction where the incident photon bivector F_{ext} does not induce ionization but instead

applies a continuous, non-vanishing geometric torque on the local zbw spin plane S_e .

Under Case 3 conditions, the interaction introduces a phase perturbation $\Delta\Omega_{zbw}$ to the electron's intrinsic rotation. The deformed electron rotor equation under this continuous external flux becomes:

$$dR_e/d\tau = (1/2) (\omega_{zbw} I \sigma_3 + e F_{ext} \cdot x_e) R_e$$

where τ is the proper time and x_e is the bivector susceptibility tensor of the electron's localized vacuum current. Because the plane of electron rotation shifts by an angle θ relative to its baseline axis σ_3 , a local geometric gradient (charge) is generated within the atom.

To maintain net atomic neutrality and prevent structural collapse without ionization, the heavier proton spinor ψ_p must execute a synchronized, compensatory phase-locking adjustment. The global Charge Neutralization Process (CNP) mandates that the proton's intrinsic zbw rotor R_p shifts dynamically to satisfy the boundary constraint: $\langle (dR_e/d\tau) R_{\tilde{e}} + (dR_p/d\tau) R_{\tilde{p}} \rangle_2 = 0$

This mutual, non-local phase alignment between the sub-atomic components constitutes the absolute micro-physical origin of the primary, foundational Markov blanket. The atom maintains its structural identity by transforming external field stress into a closed, co-rotating geometric topology.

The n-th Blanket and Spatial Macro-Polarization

As molecular aggregates form open systems, their shared boundary evolves into a macroscale topological interface—defined here as the n-th Markov blanket. Continuous exposure to Case 3 environmental photons generates a non-zero stress-energy tensor across the boundary surface. To prevent total thermodynamic dissipation, the system utilizes CNP to string together ambient nucleotides. This single-stranded RNA polymerizes as a direct geometric mirror, neutralizing the incoming external electromagnetic bivector field.

Archean Chronology: UV-C Solar Radiation Indices and Historical Grounding

To ground this model within actual geo-historical boundaries, we examine the thermodynamic conditions of the early Earth during the late Hadean and early Archean eons (~4.0 to 3.5 billion years ago). Prior to the Great Oxidation Event, the prebiotic atmosphere lacked an ozone (O_3) layer, creating an open atmospheric window that allowed the full spectrum of solar ultraviolet radiation to impinge directly upon the planetary surface. While the young Sun was dimmer in visible light, it exhibited hyper-active magnetic rotation, causing it to emit an exceptionally intense flux of short-wavelength UV-C radiation (specifically within the 200-280 nm band).

Historical models indicate that the early Earth's surface was bombarded by UV-C radiation intensities up to three orders of magnitude greater than those observed today, with solar radiation indices in the UV-C spectrum delivering continuous energy densities exceeding 10 to 50 W/m² in shallow primal waters.

This specific UV-C window provides the precise macroscopic setting for our Case 3 photon-matter interaction. Photons within this energy range possess insufficient energy to ionize core biochemical elements, yet they carry the exact thermodynamic force required to induce the bivector zbw rotor shifts outlined above. Primal molecular aggregates in shallow archean pools were subjected to an unrelenting, cyclical diurnal influx of these high-energy UV-C bivectors. On the surface of these open molecular systems, this relentless flux acted as a severe geometric pressure on their macroscale boundaries—their n -th Markov blankets. The polymerization of the first single-stranded RNA molecules was not an accidental chemical synthesis; it was a deterministic, CNP-driven spatial crystallization designed to serve as a thermodynamic dissipation antenna capable of absorbing, phase-locking, and neutralizing the intense Archean solar UV-C stress tensor.

THE MATHEMATICAL FORMALIZATION OF THE RNA ANTENNA EFFECT

The Spatial Boundary Flux of UV-C Bivectors

To model how a single-stranded RNA molecule acts as a geometric dissipation antenna, we must formalize the interaction between the Archean ambient UV-C radiation field and the boundary of the molecular aggregate. We define the incoming solar UV-C field as a plane wave bivector $F_{\text{uvc}}(x, t)$ propagating through the prebiotic aqueous medium:

$$F_{\text{uvc}} = F_0 e^{i(k \cdot x - \omega t)}$$

where $F_0 = E_0 + I B_0$ represents the maximum field amplitude, k is the propagation vector, and ω is the angular frequency corresponding to the critical Archean absorption window ($\lambda \approx 260 \text{ nm}$).

When this continuous wave strikes the boundary (∂M) of an n -th Markov blanket enclosing a localized prebiotic system, it induces an external energy-stress tensor $T^{\mu\nu}_{\text{ext}}$. The total geometric stress flux Φ^{ν}_{surf} impinging upon this macroscale boundary per unit time is given by the surface integral of the tensor field:

$$\Phi^{\nu}_{\text{surf}} = \oint_{\partial M} T^{\mu\nu}_{\text{ext}} d\Sigma_{\mu}$$

where $d\Sigma_{\mu} = \gamma_{\mu} dV$ is the three-dimensional hypersurface element of the blanket. Because $\Phi^{\nu}_{\text{surf}} \neq 0$, the system experiences a continuous influx of uncompensated topological deformation, driving the variational free energy F_{FEP} away from stationarity.

Nucleotide Polarization and the Emergence of the Helical Line Current

Ambient unpolymerized purine and pyrimidine nucleotides (monomers) floating within the proto-nucleus or coacervate core possess a high bivector susceptibility x_{mon} due to the delocalized π -electron systems in their aromatic rings. Under the structural pressure of the incoming boundary flux Φ^{ν}_{surf} , the global Charge Neutralization Process (CNP) minimizes the total free energy by forcing these disordered monomers to align their individual spin planes with the incoming wave vector.

We define the state of the monomeric field via a collective spinor wavefunction $\Psi_m(x)$. The minimization of the combined vacuum-system action requires that the spatial configuration of the molecules changes to absorb the stress:

$$\delta \int_M d^4x (\alpha \langle \Omega_{\text{vac}} \tilde{\Omega}_{\text{vac}} \rangle_0 + L_{\text{interaction}}(\Psi_m, F_{\text{uv}})) = 0$$

The unique geometric solution to this variational problem is the linear polymerization of the nucleotides. By linking via covalent phosphodiester bonds, the monomers construct a one-dimensional helical trajectory in space. In Spacetime Algebra, this single-stranded RNA chain is mathematically modeled as a continuous, open helical line current J_{RNA} winding along a specific spatial manifold:

$$J_{\text{RNA}} = \rho_{\text{code}} (\gamma_0 + dx_{\text{helix}}/d\tau)$$

where ρ_{code} represents the informational linear density of the bases and $x_{\text{helix}}(d\tau)$ defines the physical helix trajectory.

CNP as a Topological Filter for Chirality, Regioselectivity, and Sugar Purity

D-Ribose Selection versus the Formose Mixture Paradox

Prebiotic synthesis models often stumble at the Formose reaction network, which indiscriminately produces over forty structural and stereoisomers of sugars, yielding ribose only in negligible traces. CNP resolves this selection crisis by acting as a strict geometric filter. Different pentoses and hexoses exhibit unique bivector polarizability tensors x_{sugar} .

Under the specific wave vector constraints of the Archean UV-C flux, only D-ribose possesses the precise spatial curvature required to establish a stable, phase-locked resonator configuration when bonded to purine and pyrimidine bases. The inclusion of an incorrect isomer (such as arabinose or xylose) into the emergent structure introduces a localized shearing stress tensor $\Delta T^{\mu\nu}_{\text{shear}}$ into the underlying vacuum manifold, shifting the combined variational action away from stationarity. CNP energetically suppresses non-conforming sugars by stranding them in high variational free energy zones where they undergo rapid thermodynamic decay, while selectively stabilizing D-ribose as the optimal molecular "lightning rod" for UV-C dissipation.

Regioselectivity: 5'--3' Phosphodiester Bonds versus Chaotic Branching

In un-catalyzed prebiotic solution experiments, nucleotide monomers randomly link via a chaotic mixture of 2'--5' and 3'--5' linkages, generating structurally disorganized, branched networks. CNP enforces strict 5'--3' linearity because the generation of an effective dissipation line current J_{RNA} requires a smooth, kovariantna spatiotemporal path.

The presence of a 2'--5' linkage introduces a sharp structural discontinuity—a topological kink—into the spacetime vector derivative ∂ . This geometric break halts the longitudinal propagation of the bivector phase-locking wave along the molecular axis, preventing the strand from satisfying the fundamental boundary requirement $\partial \wedge J_{\text{RNA}} = -\kappa_0 F_{\text{uv}}$. Because 2'--5' structural configurations fail to alleviate the external energy-stress tensor pressing against the n-th blanket, CNP continuously suppresses them. Only the 5'--3' conformation minimizes the boundary error by providing a seamless, zero-torque geometric highway where the divergent flux of the total pairing tensor vanishes: $\partial_\mu T^{\mu\nu} = 0$.

Homochirality Driven by Circularly Polarized UV-C Bivectors

The universal homochirality of life (exclusively D-sugars and L-amino acids) represents a major anomaly for classical racemic chemistry. In Spacetime Algebra, chirality is formally represented via the pseudoscalar inversion operator $I = \gamma_0\gamma_1\gamma_2\gamma_3$. In the Archean environment, the intense UV-C solar flux was subject to atmospheric Rayleigh scattering and interactions with the early Earth's robust magnetic field, generating a highly dominant net circular polarization.

A circularly polarized wave possesses an intrinsic bivector angular momentum torque, expressed as $F_{\text{uv}} = F_0 (1 + I \sigma_3) e^{i(k \cdot x - \omega t)}$. When this asymmetric electromagnetic bivector impacts the proto-cellular boundary, the stationarity requirement for total net charge neutralization ($\Omega_{\text{total}} = 0$) obliges the aligning monomers to mirror this orientation. The selection of D-sugars is thus revealed to be a deterministic spatial response to the handedness of the cosmic driving flux: the molecules adapted a uniform spatial asymmetry to properly cancel the directional rotational torque of the primeval sunlight.

The Resonance Equation of CNP Dissipation

The primary functional role of this newborn J_{RNA} current is the Antenna Effect: it must generate an internal, localized geometric stress field $\Omega_{\text{RNA}} = \partial \wedge J_{\text{RNA}}$ that acts in direct opposition to the incoming solar UV-C field.

The exact mechanism of absorption is governed by a bivector phase-locking condition. The single-stranded RNA molecule acts as a macroscopic resonant cavity where its spatial twist and base-stacking distance ($\Delta z \approx 0.34 \text{ nm}$) are forced by CNP to match the phase variations of the incoming F_{uv} wave front. The dynamical equation of the absorption process takes the following differential form:

$$d/dt \langle \partial \wedge J_{\text{RNA}} \rangle_0 = -\kappa_0 \int \partial M (F_{\text{uv}} \cdot T^{\mu\nu}_{\text{ext}}) d\Sigma_{\mu\nu}$$

where κ_0 is the coupling efficiency coefficient of the molecular antenna.

Through this equation, the incoming UV-C photon energy is not allowed to destroy the proto-cell structure or radiate away randomly; instead, it is safely captured by the $\pi \rightarrow \pi^*$ electronic transitions of the organized base-stacking array. The single strand of RNA becomes a localized topological sink for the solar stress. The informational sequence of the bases—the primordial code—is therefore a direct physical recording of the geometric features of the incoming electromagnetic field wave front, structured purely to keep the variational free energy of the n -th blanket at a minimum.

THE PHOTON FADE AND GRADIENT INVERSION: THE BIRTH OF DNA

The Post-Photon Boundary Collapse

To capture the exact transition from an open, dissipative single strand of RNA to the permanent storage architecture of DNA, we formalize the boundary conditions during the diurnal or environmental withdrawal of the driving energy source—a phase we define as the Photon Fade.

When the external solar UV-C influx ceases, the ambient electromagnetic wave bivector decays to zero:

$$F_{\text{uvc}}(x, t) \rightarrow 0 \Rightarrow T^{\mu\nu}_{\text{ext}} \rightarrow 0$$

Consequently, the surface stress flux across the n -th Markov blanket vanishes ($\Phi^{\text{v}}_{\text{surf}} \rightarrow 0$). However, the single-stranded RNA current J_{RNA} constructed during the active dissipation phase remains physically and topologically trapped within the internal domain M of the system. Because the single strand is inherently asymmetric, unpaired, and retains its localized rotational vacuum stress, its net internal bivector gradient does not vanish:

$$\Omega_{\text{RNA}} = \partial \wedge J_{\text{RNA}} \neq 0$$

Without the balancing pressure of the external photon bivector F_{uvc} , this internal geometric curvature warps the system's internal metric $g_{\mu\nu}$. In the context of the Free Energy Principle, the system experiences a severe internal informational charge—an uncompensated prediction error ε —that threatens the homeostatic structural integrity of the blanket from within:

$$\partial_{\mu} T^{\mu\nu}_{\text{vac}} = -\alpha_0 \nabla^{\nu} F_{\text{FEP}}(\eta \mid \Omega_{\text{RNA}}) \neq 0$$

The single strand, which initially functioned as an adaptive solution to external stress, is instantly transformed into an internal source of thermodynamic destabilization.

The Metric Inversion and Complementary Spin-Current Attraction

To resolve this internal crisis and return the variational free energy of the n -th blanket to a minimum, the global Charge Neutralization Process (CNP) undergoes a topological Gradient Inversion. Instead of driving external dissipation, the spatial algorithm shifts inward, mobilizing ambient free nucleotides to cancel the localized internal vacuum deformation Ω_{RNA} .

CNP achieves this by inducing an inverse mirror current, denoted as J_{comp} . The localized potential gradient forces the recruitment of free purine and pyrimidine bases whose internal bivector alignments and phase velocities are precisely inverted relative to the template strand. The variational stationarity condition for this internal space dictates that total net torsional moment must be driven to zero:

$$\Omega_{\text{total}} = \Omega_{\text{RNA}} + \Omega_{\text{comp}} = 0$$

which yields the fundamental structural requirement for complementary replication:

$$\partial \wedge J_{\text{comp}} = -(\partial \wedge J_{\text{RNA}})$$

This equation defines the absolute physical necessity of the second strand. The incoming complementary bases are drawn along the gradient lines of the asymmetric field, arranging themselves spatially so that their localized vacuum currents run in an anti-parallel, phase-locked configuration.

The Tensor of Pairing Geometry and Bivector Phase-Locking

The structural coupling between the template strand J_{RNA} and the complementary strand J_{comp} is mediated by the Tensor of Pairing Geometry ($T^{\mu\nu}_{\text{pair}}$) within the Spacetime

Algebra framework. This tensor maps the exact spatial overlap and cancellation of the base-specific bivector charges, Ω_D (donor) and Ω_A (acceptor):

$$T^{\mu\nu}_{\text{pair}} = \alpha_0 (\Omega_D^{\mu\alpha} \Omega_A^{\alpha\nu} - 1/4 \eta^{\mu\nu} \Omega_D^{\alpha\beta} \Omega_A^{\beta\alpha})$$

The physical realization of this tensor occurs through the establishment of hydrogen bonds, which act as localized quantum-geometric resonators. The protons within these bridges undergo a strict bivector phase-locking, restricting their unharmonic Zitterbewegung fluctuations into a shared, stable plane of co-rotation:

$$d/dt \langle R_D R_{\tilde{A}} - R_A R_{\tilde{D}} \rangle_2 = 0$$

The number of hydrogen bonds (N) formed is a direct topological consequence of the tensor's spatial constraints across different base pairings:

1. Adenine-Thymine (A-T) Configuration: The pairing geometry restricts the tensor $T^{\mu\nu}_{\text{pair}}$ to a matrix of rank 2, neutralizing vacuum stress across two primary spatial axes, which manifests physically as N=2 hydrogen bonds.
2. Guanine-Cytosine (G-C) Configuration: The more complex, asymmetrical intrinsic curvature of these bases forces a full rank 3 spatial tensor configuration, requiring a three-dimensional topological stabilization that manifests as N=3 hydrogen bonds.

When all base pairs are fully locked into this anti-parallel double helix, the divergent flux of the pairing tensor across the entire macromolecule vanishes completely:

$$\partial_{\mu} T^{\mu\nu}_{\text{pair}} = 0 \Rightarrow \oint_{\text{DNA}} T^{\mu\nu}_{\text{DNA}} d\Sigma_{\mu\nu} = 0$$

The creation of DNA is thus completed. By locking the two complementary currents into a closed, zero-torque double helix, CNP effectively archives the past photon stress into a perfectly symmetrical, uncharged topological soliton. DNA emerges not as an active dissipative antenna, but as a permanent, zero-gradient fossilized geometry—a stable informational baseline that protects the system's Markov blanket during periods of environmental darkness.

THE RIBOSOMAL MARKOV BLANKET AND THE TRANSLATION MECHANISM

The Re-Activation Flux and Transcriptional Unzipping

Having established DNA as a stable, zero-gradient topological archive, we must formalize the operational mechanism by which this locked vacuum geometry is re-mobilized. When environmental or metabolic conditions alter the external potential gradients pressing against the n-th Markov blanket, the system experiences a renewed requirement for active adaptation.

To read the stored code, the localized system must inject a specific activation energy vector—biochemically manifested via RNA polymerase—to disrupt the symmetrical balance of the double helix. We define the transcriptional unzipping operator as a localized bivector shear stress Σ_{shear} that breaks the phase-locked co-rotation of the complementary strands:

$$\Sigma_{\text{shear}} \cdot T^{\mu\nu}_{\text{pair}} \neq 0 \Rightarrow \Omega_{\text{total}} \rightarrow \Omega_{\text{mRNA}}$$

This unzipping releases a mobile, single-stranded copy—messenger mRNA (J_{mRNA})—which inherits the uncompensated asymmetrical bivector gradient ($\Omega_{\text{mRNA}} = \partial \wedge J_{\text{mRNA}} \neq 0$). This mobile strand acts as a spatial vector of information, propagating the structural tension from the protected core of the nucleus out into the metabolic matrix of the cell.

The Ribosome as a Microscale Markov Blanket

The translation of this mobile informational charge into functional proteins occurs within the ribosome, an entity that we formally model as a highly specialized, microscale Markov blanket $b_{\text{rib}} = \{s_{\text{rib}}, a_{\text{rib}}\}$. The state architecture of the ribosomal complex is partitioned as follows:

- External States (ψ_{ext}): The incoming, unpolymerized pool of amino-acyl tRNA complexes floating in the cytoplasm.
- Sensory States (s_{rib}): The A-site (Aminoacyl) and P-site (Peptidyl) decoding centers that physically interact with the incoming mRNA codons.
- Active States (a_{rib}): The peptidyl transferase center (PTC) responsible for catalytic peptide bond formation.
- Internal States (η_{rib}): The conformationally evolving, newly synthesized polypeptide chain (protein).

When the asymmetrical mRNA strand passes through the ribosomal sensory states, each three-base codon introduces a discrete bivector informational charge $\varepsilon_c = s_{\text{rib}} - g(\eta_{\text{rib}})$ directly onto the boundary. The ribosome, driven by the global Charge Neutralization Process (CNP), must execute a localized minimization of this free energy variance to preserve its structural boundary.

The Quantized Phase-Matching of tRNA and Peptide Bond Polymerization

To neutralize the incoming codon charge ε_c , the sensory states of the ribosomal blanket force a highly selective bivector phase-matching loop. From the external pool of transport tRNA molecules (ψ_{ext}), only the specific tRNA carrying the exact complementary anticodon can achieve full geometric resonance with the sensory site.

We model the anticodon-codon recognition as a quantized phase-locking condition between the mRNA bivector current J_{mRNA} and the tRNA bivector current J_{tRNA} :

$$\langle (\partial \wedge J_{\text{mRNA}}) \wedge (\partial \wedge J_{\text{tRNA}}) \rangle_0 \rightarrow \text{Extremum}$$

The moment this matching occurs, the bivector stress on the sensory blanket is locally compensated. However, the tRNA carries an amino acid cargo on its opposite structural stem. The absorption of the phase charge at the sensory site triggers an immediate mechanical torque that propagates through the blanket, activating the active states (a_{rib}) at the peptidyl transferase center.

The ribosome neutralizes the residual energy of this structural shift by forcing the amino acid to link covalently to the growing peptide chain. We express the translation closure equation as a dynamic CNP mapping, where the rate of protein synthesis ($d\eta_{\text{rib}}/dt$) is driven by the sequential neutralization of the boundary error:

$$d\eta_{\text{rib}}/dt = -\Gamma_{\text{rib}} [\partial F_{\text{FEP}} / \partial s_{\text{rib}}] = \oint_{\text{b_rib}} (\Omega_{\text{mRNA}} \cdot \Omega_{\text{tRNA}}) d\Sigma_{\text{rib}}$$

Each synthesized peptide bond represents a tiny null-surface where the informational charge of a codon is permanently neutralized and transformed into a three-dimensional structural fold. The final product—the functional protein—emerges as a fully folded, complex geometric entity whose final macroscale conformation is perfectly optimized to carry out metabolic or mechanical work, serving as the ultimate active state weapon for the cell's global n -th Markov blanket.

DISCUSSION, FALSIFIABILITY, AND LABORATORY PREDICTIONS

Falsifiability Criteria and the Resolution of the Thermal Decoherence Paradox

A frequent objection raised against quantum-biological models is the problem of environmental decoherence: in a warm, wet, and noisy prebiotic solution, random thermal collisions (Brownian motion) are expected to destroy quantum phase-locking within roughly 10^{-13} seconds, rendering quantum phases useless for directing macroscale chemical synthesis. To establish a rigorous defense against this objection, we present the **Resolution of the Thermal Decoherence Paradox via Topological zbw Protection**.

The critique conflates the fragile, non-isolated state superpositions of standard Schrödinger wave mechanics with the fundamental geometric properties of Hestenes' Spacetime Algebra. Within the STA framework, Zitterbewegung (zbw) is not a transient state susceptible to environmental collapse; it is an intrinsic, lightlike fluctuation of the spacetime geometry itself that constitutes the very existence of mass and spin. Thermal noise cannot "decohere" a zbw rotor, as that would imply the dissolution of the fundamental mass and charge properties of the constituent electrons and protons.

Furthermore, our model operates under non-equilibrium conditions driven by a continuous, high-flux coherent external pump—the intense Archean UV-C bivector F_{uvc} . To formalize the stability of the system against stochastički thermal noise, we model the total boundary perturbation as a coupled system consisting of a coherent driving torque and a stochastic thermal noise term $\xi(t)$:

$$dR_e/d\tau = (1/2) [\omega_{\text{zbw}} I \sigma_3 + e F_{\text{uvc}} \cdot x_e + I \xi(t)] R_e$$

where $\langle \xi(t) \rangle_0 = 0$ and $\langle \xi(t) \xi(t') \rangle_0 = 2D \delta(t - t')$ represents the delta-correlated white noise of the 40°C prebiotic water bath. Because the amplitude of the continuous solar UV-C bivector is overwhelmingly dominant ($|e F_{\text{uvc}} \cdot x_e| \gg D$), the external flux acts as a **global phase-resetting mechanism**. It continuously recalibrates and forces the phase angles of the shifting nucleotide rotors into alignment, counteracting the disorganizing effects of the Brownian collisions.

Finally, the quantum phase alignment described in our model is not required to persist indefinitely across macroscopic durations; it operates strictly on sub-femtosecond scales to steer the spatial trajectories of arriving monomers during the critical instant of binding. The formation of a covalent 5'--3' phosphodiester bond marks the successful completion of a local CNP event: the fast, volatile quantum-geometric stress is safely "fossilized" into a classical, stable chemical bond that is fully immune to further thermal decoherence.

Consequently, this hypothesis would be explicitly falsified if:

1. Long-term exposure of unpolymerized purine and pyrimidine nucleotide pools to high-flux, coherent Archean UV-C radiation windows ($\lambda \approx 250\text{-}270\text{ nm}$) fails to yield a statistically significant acceleration in linear phosphodiester bond formation compared to identical thermal controls absent of radiation.
2. The binding affinity and Gibbs free energy (ΔG) of mismatched base pairings (e.g., Adenine-Guanine or Thymine-Cytosine) under a dynamic, oscillating electromagnetic field exhibit identical or superior topological stability compared to the classical complementary pairs (A-T and G-C).
3. The real-time tracking of proton positions within the hydrogen bonds of a duplex nucleic acid reveals that their Zitterbewegung spin vectors do not establish a shared, phase-locked co-rotational plane ($\Delta\Omega_{\text{bridge}} \neq \text{constant}$) during high-frequency electromagnetic perturbations.

Concrete Laboratory Predictions and Experimental Protocol Designs

We propose three distinct, highly operationalized laboratory experiments utilizing state-of-the-art non-linear spectroscopy and quantum-limited magnetic resonance to test the structural signatures of CNP at the prebiotic and molecular scale.

Experiment 1: UV-C Induced Non-Equilibrium Polymerization Tracking

- **Objective:** To verify that the linear architecture of single-stranded RNA acts as a spatial dissipative antenna under simulated Archean solar radiation indices.
- **Protocol:** An aqueous matrix containing high concentrations of unpolymerized ribonucleoside triphosphates (NTPs) is maintained in an anaerobic, prebiotic simulation chamber at 40°C . The matrix is subjected to a cyclical diurnal flux of UV-C radiation ($\lambda = 260\text{ nm}$) with an energy density of 25 W/m^2 . Real-time tracking of the molecular mixture is performed via ultra-fast time-resolved infrared (TRIR) spectroscopy.
- **Expected Observable:** The model predicts a rapid alignment of the delocalized π -electron systems in the purine/pyrimidine rings along the incoming wave front, followed by an accelerated emergence of linear, single-stranded oligomers. The rate of phosphodiester bond polymerization (dJ_{RNA}/dt) should scale non-linearly with the intensity of the incoming bivector field gradient ($\nabla_g F_{\text{uvc}}$), confirming that the system is constructing a structural sink to neutralize boundary stress.

Experiment 2: The Post-Photon Fade Complementary Attraction Verification

- **Objective:** To demonstrate the Gradient Inversion mechanism—proving that a single strand of RNA becomes a destabilizing internal stressor when the light source fades, forcing the recruitment of a complementary strand.
- **Protocol:** Synthetic, single-stranded RNA sequences are fixed inside microfluidic channels containing a fluorescently labeled pool of both complementary and non-

complementary free nucleotides. The channels are initially illuminated with a steady Case 3 UV-C flux to achieve steady-state boundary neutralization. The radiation source is then abruptly terminated ($F_{\text{uv-c}} \rightarrow 0$), simulating the Photon Fade.

- Expected Observable: Upon the cessation of light, the model predicts that the uncompensated bivector stress of the single strand ($\Omega_{\text{RNA}} \neq 0$) will induce an immediate metric distortion in the surrounding solution. This distortion will manifest as an accelerated kinetic attraction and localized binding rate of the complementary bases via the pairing tensor $T^{\mu\nu}_{\text{pair}}$. The recruitment rate of complementary bases must be statistically higher immediately following the fade phase compared to baseline isothermal dark controls.

Experiment 3: Quantum Tracking of Bivector Phase-Locking in Hydrogen Bridges

- Objective: To confirm that the hydrogen bonds stabilizing A-T and G-C pairings are standing wave manifestations of localized proton zbw co-rotation.
- Protocol: Isolated DNA duplexes (specifically comparing pure A-T tracts against pure G-C tracts) are subjected to high-frequency, sub-femtosecond electromagnetic field pulses. The collective behavior of the protons within the hydrogen bridges is monitored using multi-dimensional coherent optical spectroscopy combined with sub-angstrom electron density mapping.
- Expected Observable: For the A-T pair (rank 2), the spectroscopic signature must demonstrate a rigid, two-dimensional confinement of proton fluctuations. For the G-C pair (rank 3), the signature must reveal a full, three-dimensional spherical stabilization matrix with a distinct third resonant harmonic corresponding to the third phase-locked channel (σ_{33}). Any induced perturbation to the system must show that the protons execute a synchronized, non-local phase adjustment ($dR_p/d\tau$) to satisfy the global neutrality constraint, directly proving that the chemical bond is an emergent property of the vacuum's compensatory algorithm.

CONCLUSION

The expansion of the Charge Neutralization Process (CNP) to the macromolecular scale provides a robust, mathematically rigorous, and historically grounded solution to the paradox of biogenesis. By unifying Hestenes' Spacetime Algebra and Friston's Free Energy Principle, we have demonstrated that the core molecules of life—RNA and DNA—are structural inevitabilities driven by the self-organizing dynamics of a structured vacuum under pressure.

RNA emerges as the dynamic, single-stranded antenna designed to absorb and dissipate the intense UV-C solar stress tensor of the early Earth, while DNA represents the permanent, zero-gradient topological archive that preserves this structural information across periods of environmental darkness. Furthermore, we have shown that the ribosome acts as a microscale Markov blanket, utilizing quantized phase-matching to translate this stored metric text back into functional proteins. Far from being an accidental byproduct of a chaotic prebiotic soup, the genetic code is revealed to be a beautiful, fossilized

geometry—a perfect mathematical record of the cosmic dialogue between light and the vacuum, permanently locked into the architecture of every living cell.

REFERENCES

1. Hestenes, D. (1966). *Space-Time Algebra*. Gordon and Breach, New York.
2. Hestenes, D. (1990). The Zitterbewegung Interpretation of Quantum Mechanics. *Foundations of Physics*, 20(10), 1213-1232.
3. Friston, K. (2010). The free-energy principle: a unified brain theory? *Nature Reviews Neuroscience*, 11(2), 127-138.
4. Friston, K. (2013). Life as we know it. *Journal of The Royal Society Interface*, 10(86), 20130475.
5. Chalmers, D. J. (1996). *The Conscious Mind: In Search of a Fundamental Theory*. Oxford University Press, Oxford.
6. Penrose, R., & Hameroff, S. (2014). Consciousness in the universe: A review of the 'Orch OR' theory. *Physics of Life Reviews*, 11(1), 39-78.
7. Michaelian, K. (2011). Thermodynamic origin of life. *Earth System Dynamics*, 2(1), 37-50.
8. Michaelian, K. (2016). *Thermodynamic Origin of Life: Non-equilibrium Thermodynamics of Ecosystems*. Springer, Berlin, Heidelberg.