Transactions on Engineering and Computing Sciences - Vol. 12, No. 3
Publication Date: June 25, 2024
DOI:10.14738/tecs.123.17065.
Izmirli, I. M. (2024). Power Series Matrices. Transactions on Engineering and Computing Sciences, 12(3). 175-183.

SERVICES FOR SCIENCE AND EDUCATION

Power Series Matrices

Ilhan M. Izmirli
George Mason University

ABSTRACT

In this paper, we propose an algorithm that can be used to construct the elements of any *power series matrix* generated by any polynomial $p(x)$. As such, this algorithm is a generalization of the specific algorithms that give us the Pascal and the Fibonacci matrices. We will first start with the special cases of Pascal and Fibonacci matrices. The general case of power series matrices will be analyzed in Section2. We will conclude our paper by giving the algorithm in section 3.

Keywords: Pascal triangle, Fibonacci triangle, power series matrix.

INTRODUCTION

Pascal's Matrix

The binomial formula

$$(a+b)^n = \sum_{k=0}^{n} \binom{n}{k} a^{n-k} b^k$$

where the binomial coefficients $\binom{n}{k}$ can easily be computed using the addition rule

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

for any two non-negative integers n and k with $k \leq n$, is one of the most well-known formulas in elementary algebra.

It is customary to call the triangular array made up of the binomial coefficients

1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
……

the ***Pascal's triangle***. This triangle has some simple yet interesting properties that are familiar to most introductory algebra students:

1. Horizontal rows add to powers of 2, which can, of course, can easily be shown by putting $a = b = 1$ in the binomial formula.

2. The horizontal rows represent powers of 11, which can, of course, can easily be shown by putting $a = 10$ and $b = 1$ in the binomial formula.
3. Adding any two successive numbers in the diagonal containing the triangular numbers 1 , 3, 6, 10, … results in a perfect square. This, of course, is a direct consequence of the definition of triangular numbers. The n^{th} triangular number is $\frac{n(n+1)}{2}$. So the sum of two consecutive triangular numbers is
$$\frac{n(n+1)}{2} + \frac{(n+1)(n+2)}{2} = \frac{(n+1)(2n+2)}{2} = (n+1)^2$$
4. In the expansion of $(a+b)^p$, where p is a prime number, all coefficients that are greater than 1 are divisible by p; that is, if the first number to the right of 1 in any row is a prime number, then all numbers greater than 1 in that row are divisible by that prime number.

Any coefficient is of the form $\frac{p!}{(p-k)!k!}$. If p is prime, then, by definition, $(p-k)!$ and $k!$ are factors of $p!$. Therefore, since p cannot possibly be in the denominator, some multiple of p must be left in the numerator, making the coefficient an integer which is a multiple of p.

It is now acknowledged that this triangle was known well before Blaise Pascal (1623 – 1662) who "introduced" it in his famous 1653 treatise, *Traité du triangle arithmétique*. Indeed, not only the binomial coefficients, but in fact, the addition rule, which, of course, is needed to generate the coefficients, were known to Indian mathematicians[1]. For instance, according to Edwards [4], some elements of the binomial coefficients can be observed in the works of Pingala (c. 200 BC -?). A few centuries later, Varahamihira (505 CE – 587 CE) gave a clear description of the addition rule [4]2013). The triangle itself was mentioned as early as the 10th century CE, in the book *Meru-prastaara*[2] by Halayudha (?-?) See [5] for more details.

Persian mathematicians were also well acquainted with the binomial coefficients - this can be seen, for example, in the writings of Al-Karaji (953-1029) and later in those of Omar Hayyam (1048–1131), who indeed set up the entire triangle. Thus, some scholars and historians refer to the triangle as the ***Khayyam-Pascal triangle*** (See [2]).

Many other cultures were familiar with the triangle and its properties as well. For example, the triangle was known in China in the early 11th century, a fact that is, according to [9], corroborated by the works of the Jia Xian (1010–1070) and Yang Hui (1238–1298).

There were also precedents in the west. The German humanist, Petrus Apianus (1495–1552), known for his works in mathematics, astronomy, and cartography, published the full triangle in 1527. In the second half of 16th century, parts of the triangle were published by the German monk and mathematician Michael Stifel (1487–1567), and the Italian mathematicians Niccolo Fontana Tartaglia (1499–1557) and Gerolamo Cardano (1501–1576) See [2] for more details.
A closely related idea is that of the ***Pascal matrix***. The Pascal matrix is an infinite matrix containing the Pascal triangle as a submatrix. There are three convenient ways of doing this:

[1] However, keeping up with "tradition," we will, throughout this paper, refer to the triangle as Pascal's triangle.
[2] This title translates as *The Staircase of Mount Meru*.

a. As a lower triangular matrix L where the binomial coefficients are placed in rows. For example,

$$L_4 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 2 & 1 & 0 \\ 1 & 3 & 3 & 1 \end{bmatrix}$$

b. As an upper triangular matrix U where the binomial coefficients are placed in columns. For example,

$$U_4 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

c. As a symmetric matrix S where the binomial coefficients are placed on the subdiagonals. For example,

$$S_4 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \\ 1 & 3 & 6 & 10 \\ 1 & 4 & 10 & 20 \end{bmatrix}$$

See [1] for more information on Pascal matrices.

Clearly, $\det(L) = \det(U) = 1$. In fact, it can be shown that $S = LU$ and consequently,

$$\mathrm{Det}(S) = \mathrm{Det}(L)\,\mathrm{Det}(U) = 1$$

See [3] for a proof.

The Fibonacci Triangle

As is well-known, the *fibonacci sequence* $\{f_n\}$,is defined recursively as

$$f_0 = f_1 = 1$$

And for $n \geq 2$**,**

$$f_n = f_{n-1} + f_{n+2}$$

The Sequence Is Named After Leonardo Of Pisa (Fibonacci) (C.1170 – C. 1250), Who in His 1202 Book *Liber Abaci* Introduced It to The European Readers. However, As Was the Case with Pascal's Triangle, This Sequence Had Been Described Earlier by Indian Mathematicians as Well. See [6] Or [8] For More Information.

The *Fibonacci Triangle* Is A Two-Dimensional Version of The Fibonacci Sequence. It Is Defined as Follows:

$$f_{0,0} = f_{1,0} = f_{1,1} = f_{2,1} = 1$$

For $m \geq n+2$

$$f_{m,n} = f_{m-1,n} + f_{m-2,n}$$

And For $m < n+2$

$$f_{m,n} = f_{m-1,n-1} + f_{m-2,n-2}$$

So, this is a triangle with fibonacci sequences on the sides. Note that the subdiagonals are fibonacci sequences as well, except that the starting value is no longer 1. So, the left edge of the triangle (as well as the right edge) is the fibonacci sequence, the diagonal parallel to it is the fibonacci sequence, the next diagonal is the fibonacci sequence starting with $f_0 = f_1 = 2$, the next is the fibonacci sequence starting with $f_0 = f_1 = 3$, the next a fibonacci sequence starting with $f_0 = f_1 = 5$, and so on.

1
1 1
2 1 2
3 2 2 3
5 3 4 3 5
8 5 6 6 5 8
13 8 10 9 10 8 13
21 13 16 15 15 16 13 21
……

See [7] for more details.

THE CONNECTION BETWEEN THE PASCAL AND FIBONACCI MATRICES AND POWER SERIES

It Is Easy to See That If We Let

$$f(x) = 1 - x$$

For $x \neq 1$, Then the Coefficients in The Power Series Of

$$\frac{1}{[f(x)]^k}$$

Arranged in A Matrix Gives Us the Pascal Matrix. For,

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \cdots + x^\nu + \cdots$$

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \cdots + (\nu+1)x^\nu + \cdots$$

$$\frac{1}{(1-x)^3} = 1 + 3x + 6x^2 + 10x^3 + \cdots + \frac{(\nu+1)(\nu+2)}{2}x^\nu + \cdots$$

If we now form a matrix where the j^{th} row consists of the coefficients of the power series of $\frac{1}{(1-x)^{j+1}}$, for $j = 0, 1, \ldots$, we get the pascal matrix:

$$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & . & . & . \\ 1 & 2 & 3 & 4 & . & . & . & . \\ 1 & 3 & 6 & . & . & . & . & . \\ 1 & 4 & . & . & . & . & . & . \\ 1 & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . \end{bmatrix}$$

It is now natural to ask the following question: if $f(x) = 1 - x$, $x \neq 1$ is replaced by some arbitrary polynomial of degree n and the same process is applied, what types of matrices will we get?

Definition 1 *let* $p(x) = a_n x^n + \cdots a_1 x + a_0$ *with* $a_n \neq 0$ *and* $a_0 \neq 0$. *Let* $c_{j,k}$ *stand for the coefficient of* x^j *in the power series expansion of* $\frac{1}{[p(x)]^{k+1}}$, $j = 0,1,\ldots.$; $k = 0,1,\ldots$ *then, the infinite matrix* M_p

$$M_p = \begin{bmatrix} c_{0,0} & c_{0,1} & c_{0,2} & . & . & . & c_{0,k} & . & . & . \\ c_{1,0} & c_{1,1} & c_{1,2} & . & . & . & c_{1,k} & . & . & . \\ . & . & . & . & . & . & . & . & . & . \\ c_{j,0} & c_{j,1} & c_{j,2} & . & . & . & c_{j,k} & . & . & . \\ . & . & . & . & . & . & . & . & . & . \end{bmatrix}$$

Whose (j,k) *Entry Is* $c_{j,k}$ *Will Be Called the Power Series Matrix Generated By* $p(x)$.

Example 1. The simplest example is $f(x) = 1 + ax$. For $x \neq a$,

$$\frac{1}{f(x)} = \frac{1}{1+ax} = 1 - ax + a^2x^2 - a^3x^3 + a^4x^4 - a^5x^5 + \cdots + (-1)^v a^v x^v + \cdots$$

$$\frac{1}{(1+ax)^2} = -\frac{1}{a}\left(\frac{1}{f(x)}\right)' = 1 - 2ax + 3a^2x^2 - 4a^3x^3 + 5a^4x^4 - \cdots + (-1)^v(v+1)a^v x^v + \cdots$$

$$\frac{1}{(1+ax)^3} = -\frac{1}{2a}\left(\frac{1}{(1+ax)^2}\right)'$$

$$= 1 - 3ax + 6a^2x^2 - 10a^3x^3 + \cdots + (-1)^v \frac{(v+1)(v+2)}{2} a^v x^v + \cdots$$

So, the power series matrix generated by $f(x) = 1 + ax$ would be

$$\begin{bmatrix} 1 & -a & a^2 & -a^3 \ldots. \\ 1 & -2a & 3a^2 & -4a^3.. \\ 1 & -3a & 6a^2 & -10a^3. \\ \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots.. \end{bmatrix}$$

Example 2. As another example let us consider the function $p(x) = -x^2 - x + 1$. Indeed, multiplying both sides of

$$\frac{1}{p(x)} = c_0 + c_1 x + c_2 x^2 + \cdots + c_k x^k + \cdots$$

By $p(x)$, we obtain

$$1 = c_0 + (-c_0 + c_1)x + (-c_0 - c_1 + c_2)x^2 + \cdots + (-c_{k-2} - c_{k-1} + c_k)x^k + \cdots$$

Implying

$$c_0 = 1$$
$$c_1 = 1$$

And

$$c_k = c_{k-1} + c_{k-2}$$

For $k \geq 2$. Hence,

$$\frac{1}{p(x)} = 1 + x + 2x^2 + 3x^3 + 5x^4 + 8x^5 + \cdots$$

To find power series expansions of $\frac{1}{[p(x)]^k}$ we note that for any $k = 1, 2, \ldots$

$$\frac{1}{[p(x)]^{k+1}} = \frac{1}{k} \cdot \frac{1}{1+2x} \cdot \left(\frac{1}{[p(x)]^k} \right)'$$

Hence, differentiating both sides of the identity for $\frac{1}{p(x)}$ and multiplying by the power series of $\frac{1}{1-(-2x)}$ we obtain

$$\frac{1}{[p(x)]^2} = 1 + 2x + 5x^2 + 10x^3 + 20x^4 + \cdots$$

Similarly,

$$\frac{1}{[p(x)]^3} = 1 + 3x + 9x^2 + 22x^3 + \cdots$$

And so on. Consequently,

$$M_p = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 \ldots \\ 1 & 1 & 2 & 3 & 5 & 8 \ldots \\ 1 & 2 & 5 & 10 & 20 & \ldots \\ 1 & 3 & 9 & 22 & \ldots & \ldots \\ \ldots & \ldots & \ldots & \ldots & \ldots & \ldots \end{bmatrix}$$

Hence, the power series matrix generated by $p(x) = -x^2 - x + 1$, is the fibonacci matrix.

THE ALGORITHM

Now we want to give an algorithm that will give us the entries of M_p more rapidly. Let $p(x) = a_n x^n + \cdots a_1 x + a_0$ with $a_n \neq 0$ and $a_0 \neq 0$. Let $c_{j,k}$ stand for the coefficient of x^j in the power series expansion of $\frac{1}{[p(x)]^{k+1}}$, $j = 0, 1, \ldots; k = 0, 1, \ldots$

Set

$$c_{0,-1} = 1$$

And if $\mu \neq 0$, set

$$c_{\mu,-1} = 0$$

Now for $\nu = 0, 1, \ldots$
And $\mu = 0, 1, \ldots$

$$c_{\mu,\nu} = -\frac{1}{a_0}\left(a_1 c_{\mu-1,\nu} + a_2 c_{\mu-2,\nu} + \cdots + a_n c_{\mu-n,\nu} - c_{\mu,\nu-1}\right)$$

For $\mu = -n, \ldots, -1$
And $\nu = -1, 0, 1, \ldots$

$$c_{\mu,\nu} = 0$$

To see why this algorithm works, for $\lambda = -1, 0, 1, \ldots$, let us set

$$s^{(\lambda)}(x) = c_{0,\lambda} + c_{1,\lambda} x + \cdots$$

Note that the coefficient of x^j in the product

$$s^{(\lambda)}(x) p(x)$$

Is $c_{j,\lambda-1}$.
Consequently, for $\lambda = 0, 1, \ldots$

$$s^{(\lambda)}(x) p(x) = s^{(\lambda-1)}(x)$$

Since

$$s^{(-1)}(x) = c_{0,-1} + c_{1,-1} x + \cdots = 1$$

We have for $k = 0, 1, \ldots$

$$\frac{1}{(p(x))^{k+1}} = s^{(k)}(x)$$

This algorithm is, of course, a natural generalization of the addition process we apply to calculate various coefficients in pascal's triangle. In fact, in case $p(x) = 1 - x$, our algorithm simply becomes

$$c_{\mu,\nu} = c_{\mu-1,\nu} + c_{\mu,\nu-1}$$

Examples:

1. The power series matrix of $p(x) = 1 + 3x$ is

$$\begin{bmatrix} 1 & 1 & 1 & 1 & . & . & . \\ -2 & -4 & -6 & -8 & . & . & . \\ -4 & -12 & -24 & -40 & . & . & . \\ -8 & -32 & -80 & . & . & . & . \\ -16 & -80 & . & . & . & . & . \\ -32 & . & . & . & . & . & . \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & . \end{bmatrix}$$

2. The power series matrix of $p(x) = 2 - x$ is

$$\begin{bmatrix} \frac{1}{2} & \frac{1}{4} & \frac{1}{8} & \frac{1}{16} & . & . & . \\ \frac{1}{4} & \frac{2}{8} & \frac{3}{16} & \frac{4}{32} & . & . & . \\ \frac{-1}{8} & \frac{-3}{16} & \frac{-6}{32} & \frac{-10}{64} & . & . & . \\ \frac{1}{16} & \frac{4}{32} & \frac{10}{64} & . & . & . & . \\ \frac{-1}{32} & \frac{-5}{64} & . & . & . & . & . \\ \frac{1}{64} & . & . & . & . & . & . \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & . \end{bmatrix}$$

References

[1] Call, G. S. and D. J. Velleman. 1993. Pascal's matrices. American Mathematical Monthly, volume 100, (April 1993): 372–376.

[2] Coolidge, J. L. 1949. The story of the binomial theorem. The American Mathematical Monthly 56: 147–157.

[3] Edelman Alan, and Gilbert Strang. 2004. Pascal Matrices. American Mathematical Monthly 111 (3): 361–385

[4] Edwards, A. W. F. 2002. Pascal's arithmetical triangle: the story of a mathematical idea. John Hopkins University Press.

[5] Edwards, A. W. F. 2013. The arithmetical triangle. In Wilson, Robin; Watkins, John J., Combinatorics: Ancient and Modern, Oxford University Press, pp. 166–180.

[6] Goonatilake, Susantha. 1998, Toward a Global Sciece. Indiana University Press,

[7] Hoggartt Jr., V. E. and Marjorie Bricknell. 1974. Triangular Numbers. The Fibonacci Quarterly. Vol 12, No 3, Ocy 1974, pp 221 – 230.

[8] Siegler, Laurence E. 2002, Fibonacci's Liber Abaci: A Translation into Modern English of the Book of Calculation, Sources and Studies in the History of Mathematics and Physical Sciences. Springer.

[9] Smith, Karl J. 2010, Nature of Mathematics, Cengage Learning,